

EXT VANISHING AND INFINITE AUSLANDER-BUCHSBAUM

PETER JØRGENSEN

ABSTRACT. A vanishing theorem is proved for Ext groups over non-commutative graded algebras. Along the way, an “infinite” version is proved of the non-commutative Auslander-Buchsbaum theorem.

0. INTRODUCTION

Let R be a noetherian local commutative ring, and let X be a finitely generated R -module of finite projective dimension. The classical Auslander-Buchsbaum theorem states

$$\mathrm{pd} X = \mathrm{depth} R - \mathrm{depth} X.$$

This can also be phrased as an Ext vanishing theorem, namely, if M is any R -module, then

$$\mathrm{Ext}_R^i(X, M) = 0 \quad \text{for } i > \mathrm{depth} R - \mathrm{depth} X. \quad (1)$$

In [1] is proved a surprising variation of this: Suppose that R is complete in the $\mathfrak{m}$ -adic topology. Then equation (1) remains true if X is *any* R -module of finite projective dimension, provided M is finitely generated. In other words, the condition of being finitely generated is shifted from X to M .

In theorem 2.4 below, this result will be generalized to the situation of a non-commutative noetherian $\mathbb{N}$ -graded connected algebra.

The route goes through an “infinite” version of the non-commutative Auslander-Buchsbaum theorem, given in theorem 1.4. This result is a substantial improvement of the original non-commutative Auslander-Buchsbaum theorem, as given in [3, thm. 3.2], in that the condition of dealing only with finitely generated modules is dropped.

The notation of this paper is standard, and is already on record in several places such as [2] or [3]. So I will not say much, except that throughout, k is a field, and A is a noetherian $\mathbb{N}$ -graded connected

2000 *Mathematics Subject Classification.* 16E30, 16W50.

Key words and phrases. k -flat dimension, depth, infinite non-commutative Auslander-Buchsbaum theorem, Ext groups, vanishing theorem.

k -algebra. However, let me give one important word of caution: Everything in sight is graded. So for instance, $D(A)$ stands for $D(\text{Gr } A)$, the derived category of the abelian category $\text{Gr}(A)$ of $\mathbb{Z}$ -graded A -left-modules and graded homomorphisms of degree zero.

1. AUSLANDER-BUCHSBAUM

Definition 1.1. For X in $D(A)$, define the k -flat dimension by

$$\text{k.fd } X = -\inf k \overset{\text{L}}{\otimes}_A X.$$

Remark 1.2. Using a minimal free resolution, it is easy to see that if the cohomology of X is bounded and finitely generated, then

$$\text{k.fd } X = \text{fd } X = \text{pd } X,$$

where fd stands for flat dimension and pd stands for projective dimension.

In the following lemma is used $D^b(A)$, the full subcategory of $D(A)$ consisting of complexes with bounded cohomology, and $D^+(A^{\text{op}})$, the full subcategory of $D(A^{\text{op}})$ consisting of complexes with cohomology vanishing in low degrees.

Lemma 1.3. *Let X in $D^b(A)$ have $\text{fd } X < \infty$, and let S in $D^+(A^{\text{op}})$ have $\dim_k h^i(S) < \infty$ for each i . Then*

$$\inf S \overset{\text{L}}{\otimes}_A X = \inf S - \text{k.fd } X.$$

Proof. Observe that $\text{fd } X < \infty$ implies

$$\inf k \overset{\text{L}}{\otimes}_A X > -\infty,$$

hence

$$\text{k.fd } X < \infty. \tag{2}$$

If

$$\inf S = \infty \tag{3}$$

then S is zero. Then $S \overset{\text{L}}{\otimes}_A X$ is also zero, so

$$\inf S \overset{\text{L}}{\otimes}_A X = \infty. \tag{4}$$

Combining equations (2), (3), and (4) shows that for $\inf S = \infty$, the lemma's equation reads

$$\infty = \infty - (\text{something smaller than } \infty)$$

which is true.

So for the rest of the proof, I can assume $\inf S < \infty$, that is, S is non-zero. Note that S is in $D^+(A^{\text{op}})$, so $\inf S > -\infty$, so $\inf S$ is a finite number.

Let me first do the case where S is concentrated in degree zero. Here S is just a graded A -right-module which is finite dimensional over k , and the lemma claims

$$\inf S \overset{\text{L}}{\otimes}_A X = -\text{k.fd } X = \inf k \overset{\text{L}}{\otimes}_A X. \quad (5)$$

To prove this, note that $\dim_k S < \infty$ gives that there is a short exact sequence $0 \rightarrow k(\ell) \rightarrow S \rightarrow \tilde{S} \rightarrow 0$ of graded A -right-modules. This gives a distinguished triangle $k(\ell) \rightarrow S \rightarrow \tilde{S} \rightarrow$ in $D(A^{\text{op}})$, and tensoring with X and taking the cohomology long exact sequence gives a sequence consisting of pieces

$$h^i(k(\ell) \overset{\text{L}}{\otimes}_A X) \rightarrow h^i(S \overset{\text{L}}{\otimes}_A X) \rightarrow h^i(\tilde{S} \overset{\text{L}}{\otimes}_A X).$$

Induction on $\dim_k S$ now shows that the lowest i with

$$h^i(k \overset{\text{L}}{\otimes}_A X) \neq 0$$

equals the lowest i with

$$h^i(S \overset{\text{L}}{\otimes}_A X) \neq 0,$$

and equation (5) follows. The induction works because $\inf k \overset{\text{L}}{\otimes}_A X > -\infty$, and even works for $\inf k \overset{\text{L}}{\otimes}_A X = \infty$.

Let me next do the case where S is general. There is a standard spectral sequence

$$E_2^{pq} = h^p(h^q(S) \overset{\text{L}}{\otimes}_A X) \Rightarrow h^{p+q}(S \overset{\text{L}}{\otimes}_A X).$$

Since $\text{fd } X < \infty$, the spectral sequence is first quadrant up to shift, hence converges. Now, $\dim_k h^q(S) < \infty$ holds for each q , so if $h^q(S)$ is non-zero, then the special case of the lemma dealt with above applies to $h^q(S) \overset{\text{L}}{\otimes}_A X$ and shows

$$\inf h^q(S) \overset{\text{L}}{\otimes}_A X = -\text{k.fd } X. \quad (6)$$

There are now two cases: First, if

$$\text{k.fd } X = -\infty \quad (7)$$

then equation (6) gives that if $h^q(S)$ is non-zero, then $\inf h^q(S) \overset{\text{L}}{\otimes}_A X = \infty$, that is, $h^p(h^q(S) \overset{\text{L}}{\otimes}_A X)$ is zero for each p . Of course this also holds for $h^q(S)$ equal to zero, and so in the spectral sequence, E_2^{pq}

is identically zero. Therefore the limit $\mathrm{h}^{p+q}(S \overset{\mathrm{L}}{\otimes}_A X)$ of the spectral sequence is also zero, so $S \overset{\mathrm{L}}{\otimes}_A X$ is zero, so

$$\inf S \overset{\mathrm{L}}{\otimes}_A X = \infty. \quad (8)$$

But $\inf S$ is a finite number, and combining this with equations (7) and (8) says that the lemma's equation reads

$$\infty = (\text{a finite number}) - (-\infty)$$

which is true.

Secondly, if

$$\mathrm{k.f.d} X > -\infty,$$

then equation (6) says that if $\mathrm{h}^q(S)$ is non-zero, then $\mathrm{h}^p(\mathrm{h}^q(S) \overset{\mathrm{L}}{\otimes}_A M)$ is non-zero for $p = -\mathrm{k.f.d} X$, but zero for $p < -\mathrm{k.f.d} X$. And of course, if $\mathrm{h}^q(S)$ is zero, then $\mathrm{h}^p(\mathrm{h}^q(S) \overset{\mathrm{L}}{\otimes}_A X)$ is zero for each p . So in the spectral sequence, E_2^{pq} is non-zero for $p = -\mathrm{k.f.d} X$ and $q = \inf S$, but zero for lower p or q . Hence $E_2^{-\mathrm{k.f.d} X, \inf S}$ can be used in a standard corner argument which shows that the lowest non-zero term in the limit $\mathrm{h}^{p+q}(S \overset{\mathrm{L}}{\otimes}_A X)$ of the spectral sequence has degree $p + q = -\mathrm{k.f.d} X + \inf S$. Hence

$$\inf S \overset{\mathrm{L}}{\otimes}_A X = -\mathrm{k.f.d} X + \inf S,$$

proving the lemma's equation. $\square$

Theorem 1.4 (Infinite Auslander-Buchsbaum). *Assume that A satisfies $\dim_k \mathrm{Ext}_A^i(k, A) < \infty$ for each i . Let X in $\mathrm{D}^b(A)$ have $\mathrm{fd} X < \infty$. Then*

$$\mathrm{depth} X = \mathrm{depth} A - \mathrm{k.f.d} X.$$

Proof. I have

$$\mathrm{RHom}_A(k, X) \cong \mathrm{RHom}_A(k, A \overset{\mathrm{L}}{\otimes}_A X) \cong \mathrm{RHom}_A(k, A) \overset{\mathrm{L}}{\otimes}_A X,$$

where the second $\cong$ holds because it holds if X is just a graded flat module, so also holds if X is quasi-isomorphic to a bounded complex of graded flat modules, and this is the case since X is in $\mathrm{D}^b(A)$ and has $\mathrm{fd} X < \infty$.

Thus

$$\begin{aligned}
\text{depth } X &= \inf \text{RHom}_A(k, X) \\
&= \inf \text{RHom}_A(k, A) \overset{\text{L}}{\otimes}_A X \\
&\stackrel{(a)}{=} \inf \text{RHom}_A(k, A) - \text{k.fd } X \\
&= \text{depth } A - \text{k.fd } X,
\end{aligned}$$

where (a) is by lemma 1.3. The lemma applies because $\text{RHom}_A(k, A)$ is in $\mathbf{D}^+(A^{\text{op}})$, and has $\dim_k \text{h}^i \text{RHom}_A(k, A) = \dim_k \text{Ext}_A^i(k, A) < \infty$ for each i by assumption. $\square$

Remark 1.5. Theorem 1.4 even holds for $\text{depth } A = \infty$, where the theorem states $\text{depth } X = \infty$.

On the other hand, suppose $\text{depth } A < \infty$. Then it is easy to see that it makes sense to rearrange the equation in theorem 1.4 as

$$\text{k.fd } X = \text{depth } A - \text{depth } X.$$

If the cohomology of X is bounded and finitely generated, then this equation reads

$$\text{pd } X = \text{depth } A - \text{depth } X$$

by remark 1.2. This is the original non-commutative Auslander-Buchsbaum theorem, as proved in [3, thm. 3.2].

2. EXT VANISHING

Lemma 2.1. *Let X in $\mathbf{D}^b(A)$ have $\text{fd } X < \infty$, and let T in $\mathbf{D}^+(A^{\text{op}})$ be so that $\text{h}^i(T)$ is a graded torsion module for each i . Then*

$$\inf T \overset{\text{L}}{\otimes}_A X \geq \inf T - \text{k.fd } X.$$

Proof. As in the proof of lemma 1.3, it is easy to see that for $\inf T = \infty$, the lemma's inequality trivially reads $\infty \geq \infty$. So for the rest of the proof I can assume $\inf T < \infty$. Note that T is in $\mathbf{D}^+(A^{\text{op}})$, so $\inf T > -\infty$, so $\inf T$ is a finite number.

Let me first do the case where T is concentrated in degree zero. Here T is just a graded torsion module, and the lemma claims

$$\inf T \overset{\text{L}}{\otimes}_A X \geq -\text{k.fd } X = \inf k \overset{\text{L}}{\otimes}_A X.$$

If F is a flat resolution of X , then this amounts to

$$\inf T \otimes_A F \geq \inf k \otimes_A F. \tag{9}$$

To prove this, note that as T is a graded torsion module, it is the colimit of the system

$$T\langle 1 \rangle \subseteq T\langle 2 \rangle \subseteq \cdots$$

where

$$T\langle j \rangle = \{ t \in T \mid A_{\geq j} t = 0 \}.$$

Each quotient $T\langle j \rangle / T\langle j-1 \rangle$ is annihilated by $A_{\geq 1}$ so has the form $\coprod_{\alpha} k(\ell_{\alpha})$, so there are short exact sequences of the form

$$0 \rightarrow T\langle j-1 \rangle \longrightarrow T\langle j \rangle \longrightarrow \coprod_{\alpha} k(\ell_{\alpha}) \rightarrow 0.$$

Tensoring such a sequence with F gives a short exact sequence of complexes because F consists of graded flat modules. The corresponding cohomology long exact sequence consists of pieces

$$h^i(T\langle j-1 \rangle \otimes_A F) \longrightarrow h^i(T\langle j \rangle \otimes_A F) \longrightarrow \coprod_{\alpha} h^i(k \otimes_A F)(\ell_{\alpha}).$$

Induction on j now makes clear that

$$h^i(k \otimes_A F) = 0$$

implies

$$h^i(T\langle j \rangle \otimes_A F) = 0 \text{ for each } j,$$

and this further gives

$$h^i(T \otimes_A F) \cong h^i(\operatorname{colim} T\langle j \rangle \otimes_A F) \cong \operatorname{colim} h^i(T\langle j \rangle \otimes_A F) = 0,$$

so the inequality (9) follows.

Let me next do the case where T is general. There is a standard spectral sequence

$$E_2^{pq} = h^p(h^q(T) \overset{L}{\otimes}_A X) \Rightarrow h^{p+q}(T \overset{L}{\otimes}_A X).$$

Since $\operatorname{fd} X < \infty$, the spectral sequence is first quadrant up to shift, hence converges. Now, each $h^q(T)$ is a graded torsion module, so if $h^q(T)$ is non-zero, then the special case of the lemma dealt with above applies to $h^q(T) \overset{L}{\otimes}_A X$ and shows

$$\inf h^q(T) \overset{L}{\otimes}_A X \geq -\operatorname{k.f.d.} X.$$

Hence if $h^q(T)$ is non-zero, then $h^p(h^q(T) \overset{L}{\otimes}_A X)$ is zero for $p < -\operatorname{k.f.d.} X$. And of course, if $h^q(T)$ is zero, then $h^p(h^q(T) \overset{L}{\otimes}_A X)$ is zero for each p . So in the spectral sequence, E_2^{pq} is zero for $p < -\operatorname{k.f.d.} X$ or $q < \inf T$. Therefore the limit $h^{p+q}(T \overset{L}{\otimes}_A X)$ of the spectral sequence is zero for $p+q < -\operatorname{k.f.d.} X + \inf T$, and this says

$$\inf T \overset{L}{\otimes}_A X \geq -\operatorname{k.f.d.} X + \inf T,$$

proving the lemma's inequality. $\square$

Lemma 2.2. *Let A satisfy $\text{depth } A < \infty$ and $\dim_k \text{Ext}_A^i(k, A) < \infty$ for each i .*

Let X in $\mathbf{D}^b(A)$ have $\text{fd } X < \infty$, and let T in $\mathbf{D}^+(A^{\text{op}})$ be so that $h^i(T)$ is a graded torsion module for each i . Then

$$\inf T \overset{\text{L}}{\otimes}_A X \geq \inf T + \text{depth } X - \text{depth } A.$$

Proof. Using lemma 2.1 and remark 1.5 gives $\geq$ and $=$ in

$$\inf T \overset{\text{L}}{\otimes}_A X \geq \inf T - \text{k.fd } X = \inf T + \text{depth } X - \text{depth } A.$$

□

In the following theorem is used $\mathbf{D}_{\text{fg}}^-(A)$, the full subcategory of $\mathbf{D}(A)$ consisting of complexes whose cohomology vanishes in high degrees and consists of finitely generated graded modules.

Theorem 2.3. *Let A satisfy $\text{depth } A < \infty$ and $\dim_k \text{Ext}_A^i(k, A) < \infty$ for each i .*

Let X in $\mathbf{D}^b(A)$ have $\text{fd } X < \infty$, and let M be in $\mathbf{D}_{\text{fg}}^-(A)$. Then

$$\sup \text{RHom}_A(X, M) \leq \sup M - \text{depth } X + \text{depth } A.$$

Proof. It is easy to see that since M is in $\mathbf{D}_{\text{fg}}^-(A)$, the Matlis dual M' is in $\mathbf{D}^+(A^{\text{op}})$ and has $h^i(M')$ a graded torsion module for each i . So

$$\begin{aligned} \sup \text{RHom}_A(X, M) &= \sup \text{RHom}_A(X, M'') \\ &\stackrel{(a)}{=} \sup((M' \overset{\text{L}}{\otimes}_A X)') \\ &= -\inf M' \overset{\text{L}}{\otimes}_A X \\ &\stackrel{(b)}{\leq} -\inf M' - \text{depth } X + \text{depth } A \\ &= \sup M - \text{depth } X + \text{depth } A, \end{aligned}$$

where (a) is by adjunction and (b) is by lemma 2.2. □

The following is the special case of theorem 2.3 where X and M are concentrated in degree zero, that is, where X and M are graded modules.

Theorem 2.4 (Ext vanishing). *Assume that A satisfies $\text{depth } A < \infty$ and $\dim_k \text{Ext}_A^i(k, A) < \infty$ for each i .*

Let X in $\text{Gr}(A)$ have $\text{fd } X < \infty$, and let M be in $\text{gr}(A)$. Then

$$\text{Ext}_A^i(X, M) = 0 \text{ for } i > \text{depth } A - \text{depth } X.$$

This says that for $\text{fd } X < \infty$, the number $\text{depth } A - \text{depth } X$ plays the role of projective dimension of X , but only with respect to finitely generated graded modules M .

Of course, this fails when M is general, as illustrated by the following example.

Example 2.5. Let A be the polynomial algebra $k[x]$. Then the conditions of theorem 2.4 are satisfied, and it is classical that $\text{depth } A$ is 1.

Let X be $k[x, x^{-1}]$. Then $\text{depth } X \geq 1$ because X is a graded torsion free module, so $\text{depth } A - \text{depth } X \leq 0$ and theorem 2.4 gives

$$\text{Ext}_A^i(X, M) = 0 \quad \text{for } i > 0$$

for M in $\text{gr}(A)$.

However, this must fail when M is general, for otherwise X would be a projective object of $\text{Gr}(A)$ which it is certainly not.

Acknowledgement. I thank Anders Frankild for showing me [1].

REFERENCES

- [1] A. Frankild, *Vanishing of local homology*, to appear in Math. Z.
- [2] P. Jørgensen, *Gorenstein homomorphisms of non-commutative rings*, J. Algebra **211** (1999), 240–267.
- [3] P. Jørgensen, *Non-commutative graded homological identities*, J. London Math. Soc. (2) **57** (1998), 336–350.

DANISH NATIONAL LIBRARY OF SCIENCE AND MEDICINE, NØRRE ALLÉ 49,
2200 KØBENHAVN N, DK-DENMARK

E-mail address: pej@dnlib.dk, www.geocities.com/popjoerg